

ERRATUM

Open Access



Erratum to: 'Perioperative use of crystalloids in patients undergoing open radical cystectomy: balanced Ringer's maleate versus a glucose 5 %/potassium-based balanced solution: study protocol for a randomized controlled trial'

Lukas M. Löffel¹, Bettina Kleeb¹, Fiona C. Burkhard² and Patrick Y. Wuethrich^{1*}

Unfortunately, the original version of this article [1] contained a few errors in Table 1. The correct data for Table 1 has now been included below.

Author details

¹Department of Anesthesiology and Pain Medicine, Inselspital, Bern University Hospital, Berne CH-3010, Switzerland. ²Department of Urology, Inselspital, Bern University Hospital, Bern, Switzerland.

Published online: 18 February 2016

Reference

1. Löffel LM, Bettina K, Burkhard FC, Wuethrich PY. Perioperative use of crystalloids in patients undergoing open radical cystectomy: balanced Ringer's maleate versus a glucose 5 %/potassium-based balanced solution: study protocol for a randomized controlled trial. *Trials*. 2014;15:276.

* Correspondence: patrick.wuethrich@insel.ch

¹Department of Anesthesiology and Pain Medicine, Inselspital, Bern University Hospital, Berne CH-3010, Switzerland

Table 1 Electrolyte composition of the 2 different crystalloid solutions

	Ringerfundin®	G5-K Solution®
Natrium	145.0 mmol/l	50.0 mmol/l
Kalium	4.0 mmol/l	30.0 mmol/l
Magnesium	1.0 mmol/l	2.0 mmol/l
Calcium	2.5 mmol/l	0 mmol/l
Chlorid	127.0 mmol/l	65 mmol/l
Acetat	24.0 mmol/l	0 mmol/l
Maleat acid	5.0 mmol/l	0 mmol/l
Lactate	0 mmol/l	18 mmol/l
HPO ₄	0 mmol/l	8.0 mmol/l
Glucose	0 g/l	50 g/l