

CORRECTION

Open Access



# Correction: Bipolar haemostatic forceps versus standard therapy by haemoclip + / – epinephrine injection as initial endoscopic treatment in active non-variceal upper GI bleeding: study protocol for a prospective, randomized multicentre trial (BeBop-Trial)

Daniel Schmitz<sup>1\*</sup>, Lucas Thielemann<sup>1</sup> and Felix Grassmann<sup>2</sup>

**Correction:** *Trials* 24, 407 (2023)

<https://doi.org/10.1186/s13063-023-07394-x>

Following publication of the original article [1], we have been notified that the financial supporter of the study has changed. Therefore, Additional files 2 and 3 have been updated, as well as the Funding section.

Originally published Funding section:

Open Access funding enabled and organized by Projekt DEAL. Financial support for the study is provided by PENTAX Medical Europe, Julius-Vosselerstraße 104, 22527 Hamburg, Germany. Specifically, an expense allowance of 75 € will be paid for each patient recruited. In addition, the costs for applying for the ethical vote will

be covered by Pentax. No further funding will be provided. The funder has had and will have no influence on the study design, data collection, data management, data analysis, interpretation of the data, writing of the report, and the decision to submit the report for publication (Additional files 2 and 3).

Corrected Funding section:

Open Access funding enabled and organized by Projekt DEAL. Financial support for the study is provided by Helios Kliniken GmbH, Germany (Grant no. 2022-0035). Specifically, an expense allowance of 75 € will be paid for each patient recruited. In addition, the costs for applying for the ethical vote will be covered by Helios Kliniken GmbH, Germany. No further funding will be provided. The funder has, had, and will have no influence on the study design, data collection, data management, data analysis, interpretation of the data, writing of the report, and the decision to submit the report for publication (Additional files 2 and 3).

The original article has been corrected.

The original article can be found online at <https://doi.org/10.1186/s13063-023-07394-x>.

\*Correspondence:

Daniel Schmitz  
[Daniel.Schmitz@helios-gesundheit.de](mailto:Daniel.Schmitz@helios-gesundheit.de)

<sup>1</sup> Department of Gastroenterology and Infectiology, Helios Kliniken Schwerin, University Campus of Medical School Hamburg, Wismarsche Str.393-397, Schwerin 19055, Germany

<sup>2</sup> Department of Medical Statistics and Epidemiology, Medical School Hamburg, Am Kaiserkai 1, Hamburg 20457, Germany



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13063-023-07751-w>.

**Additional file 2.** Contract funder/coordinating study centre (German) from 31 December 2022.

**Additional file 3.** Contract funder/coordinating study centre (translated in English) from 31 December 2022.

Published online: 10 November 2023

## Reference

1. Schmitz D, et al. Bipolar haemostatic forceps versus standard therapy by haemoclip + / – epinephrine injection as initial endoscopic treatment in active non-variceal upper GI bleeding: study protocol for a prospective, randomized multicentre trial (BeBop-Trial). *Trials*. 2023;24:407. <https://doi.org/10.1186/s13063-023-07394-x>.