

POSTER PRESENTATION

Open Access

Using re-randomisation to improve patient recruitment and increase statistical power in clinical trials

Brennan Kahan^{1*}, Andrew Forbes², Caroline Dore³, Tim Morris⁴

From 3rd International Clinical Trials Methodology Conference
Glasgow, UK. 16-17 November 2015

Patient recruitment is a major challenge for randomised trials. Reviews of publicly funded UK trials have found that 45 to 69% fail to recruit to target. This increases costs, delays results, and adversely impacts on the feasibility of conducting trials for conditions where there is a limited patient pool.

For some conditions a patient requires treatment on multiple occasions. For example, patients with sickle cell disease require pain relief for each pain crisis, or women having fertility treatment undergo multiple treatment cycles until becoming pregnant. The current norm is for patients to be included for only one treatment episode. We propose a re-randomisation design, allowing patients to be randomised on multiple occasions, which could substantially increase the recruitment rate.

We describe some properties of the re-randomisation design, such as the conditions required to obtain unbiased estimates of the treatment effect and control type I error rates, and offer advice on practical design and analysis issues. We show that this design can reduce the required number of patients compared with a parallel group design. It can be used in a wider variety of settings than a cross-over design and may be more patient-centred as the number of treatment periods is determined by participants rather than the trial design. We also highlight situations where this design would be inappropriate.

Authors' details

¹Pragmatic Clinical Trials Unit, Queen Mary University of London, London, UK. ²School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia. ³Comprehensive Clinical Trials Unit, University College London, London, UK. ⁴Hub for Trials Methodology Research, MRC Clinical Trials Unit at UCL, London, UK.

¹Pragmatic Clinical Trials Unit, Queen Mary University of London, London, UK
Full list of author information is available at the end of the article

Published: 16 November 2015

doi:10.1186/1745-6215-16-S2-P219

Cite this article as: Kahan *et al.*: Using re-randomisation to improve patient recruitment and increase statistical power in clinical trials. *Trials* 2015 **16**(Suppl 2):P219.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

