

CORRECTION

Open Access



Correction to: Comparison of the effectiveness of a tailored cognitive behavioural therapy with a supportive listening intervention for depression in those newly diagnosed with multiple sclerosis (the ACTION-MS trial): protocol of an assessor-blinded, active comparator, randomised controlled trial

Litza Kiropoulos^{1,2*}, Trevor Kilpatrick^{2,3}, Tomas Kalincik², Leonid Churilov³, Elizabeth McDonald⁴, Tissa Wijeratne⁵, Jennifer Threader¹, Vanja Rozenblat¹, Neil O'Brien-Simpson⁶, Anneke Van Der Walt⁷ and Lisa Taylor²

Correction to: *Trials* (2020) 21: 100
<https://doi.org/10.1186/s13063-019-4018-8>

After publication of our article [1] the authors have notified us that three of the names have been incorrectly spelled.

- Original name spelling:
Tomas Kalincek
Leonid Cherulov
Neil Simpson-O'Brien
- Correct name spelling:
Tomas Kalincik
Leonid Churilov
Neil O'Brien-Simpson

Author details

¹Melbourne School of Psychological Sciences, University of Melbourne, Melbourne, Victoria 3010, Australia. ²Department of Neurology, Royal Melbourne Hospital, Melbourne, Victoria, Australia. ³Florey Institute of Neuroscience and Mental Health, Melbourne, Victoria, Australia. ⁴Department of Neuroscience, Rehabilitation and Neuroimmunology, St Vincent's Hospital, Melbourne, Victoria, Australia. ⁵Department of Neurology, Western Health, Sunshine, Victoria, Australia. ⁶Melbourne Dental School, University of Melbourne, Carlton, Victoria, Australia. ⁷Department of Neurosciences, Monash University, Melbourne, Victoria, Australia.

Published online: 20 March 2020

Reference

1. Kiropoulos L, et al. Comparison of the effectiveness of a tailored cognitive behavioural therapy with a supportive listening intervention for depression in those newly diagnosed with multiple sclerosis (the ACTION-MS trial): protocol of an assessor-blinded, active comparator, randomised controlled trial. *Trials*. 2020;21:100. <https://doi.org/10.1186/s13063-019-4018-8>.

The original article can be found online at <https://doi.org/10.1186/s13063-019-4018-8>

* Correspondence: litzak@unimelb.edu.au

¹Melbourne School of Psychological Sciences, University of Melbourne, Melbourne, Victoria 3010, Australia

²Department of Neurology, Royal Melbourne Hospital, Melbourne, Victoria, Australia

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.