

LETTER

Open Access



Proactive Prophylaxis With Azithromycin and HydroxyChloroquine in Hospitalised Patients With COVID-19 (ProPAC-COVID): A structured summary of a study protocol for a randomised controlled trial

Pradeesh Sivapalan^{1,2}, Charlotte Suppli Ulrik³, Rasmus Dahlin Bojesen⁴, Therese Sophie Lapperre⁵, Josefin Viktoria Eklöf¹, Kjell Erik Julius Håkansson³, Andrea Browatzki⁶, Casper Tidemansen³, Jon Torgny Wilcke¹, Julie Janner³, Vibeke Gottlieb¹, Howraman Meteran¹, Celeste Porsbjerg⁵, Birgitte Lindegaard Madsen⁶, Mia Moberg³, Lars Pedersen⁵, Thomas Lars Benfield⁷, Jens Dilling Lundgren⁸, Filip Krag Knop⁹, Tor Biering-Sørensen¹⁰, Muzhda Ghanizada⁵, Tine Peick Sonne¹, Uffe Christian Steinholtz Bødtger^{2,11,12}, Sidse Graff Jensen¹, Daniel Bech Rasmussen¹¹, Eva Brøndum³, Oliver Djurhuus Tupper³, Susanne Wiemann Sørensen³, Gitte Alstrup¹¹, Christian Borbjerg Laursen¹³, Ulla Weinrich Møller¹⁴, Asger Sværrild⁵ and Jens-Ulrik Stæhr Jensen^{1*} 

Abstract

Objectives: The aim of this randomised GCP-controlled trial is to clarify whether combination therapy with the antibiotic azithromycin and hydroxychloroquine via anti-inflammation/immune modulation, antiviral efficacy and pre-emptive treatment of supra-infections can shorten hospitalisation duration for patients with COVID-19 (measured as "days alive and out of hospital" as the primary outcome), reduce the risk of non-invasive ventilation, treatment in the intensive care unit and death.

Trial design: This is a multi-centre, randomised, Placebo-controlled, 2-arm ratio 1:1, parallel group double-blind study.

Participants: 226 participants are recruited at the trial sites/hospitals, where the study will take place in Denmark: Aalborg, Bispebjerg, Gentofte, Herlev, Hillerød, Hvidovre, Odense and Slagelse hospitals.

Inclusion criteria:

- Patient admitted to Danish emergency departments, respiratory medicine departments or internal medicine departments
- Age \geq 18 years

(Continued on next page)

* Correspondence: jens.ulrik.jensen@regionh.dk

¹Section of Respiratory Medicine, Department of Medicine, Herlev and Gentofte Hospital University of Copenhagen, Hellerup, Denmark
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

- Hospitalized ≤ 48 hours
- Positive COVID-19 test / diagnosis during the hospitalization (confirmed).
- Men or non-fertile women. Fertile women* must not be pregnant, i.e. negative pregnancy test must be available at inclusion
- Informed consent signed by the patient

*Defined as after menarche and until postmenopausal (no menstruation for 12 months)

Exclusion criteria:

- At the time of recruitment, the patient uses >5 LO₂/min (equivalent to 40% FiO₂ if measured)
- Known intolerance/allergy to azithromycin or hydroxychloroquine or hypersensitivity to quinine or 4-aminoquinoline derivatives
- Neurogenic hearing loss
- Psoriasis
- Retinopathy
- Maculopathy
- Visual field changes
- Breastfeeding
- Severe liver diseases other than amoebiasis (INR > 1.5 spontaneously)
- Severe gastrointestinal, neurological and hematological disorders (investigator-assessed)
- eGFR < 45 ml/min/1.73 m²
- Clinically significant cardiac conduction disorders/arrhythmias or prolonged QTc interval (QTc (f) of $> 480/470$ ms).
- Myasthenia gravis
- Treatment with digoxin*
- Glucose-6-phosphate dehydrogenase deficiency
- Porphyria
- Hypoglycaemia (Blood glucose at any time since hospitalization of < 3.0 mmol/L)
- Severe mental illness which significantly impedes cooperation
- Severe linguistic problems that significantly hinder cooperation
- Treatment with ergot alkaloids

*The patient must not be treated with digoxin for the duration of the intervention. For atrial fibrillation/flutter, select according to the Cardiovascular National Treatment Guide (NBV): Calcium antagonist, Beta blocker, direct current (DC) conversion or amiodarone. In case of urgent need for digoxin treatment (contraindication for the aforementioned equal alternatives), the test drug should be paused, and ECG should be taken daily.

Intervention and comparator: Control group:

The control group will receive the standard treatment + placebo for both types of intervention medication at all times. If part or all the intervention therapy being investigated becomes standard treatment during the study, this may also be offered to the control group.

Intervention group:

The patients in the intervention group will also receive standard care. Immediately after randomisation to the intervention group, the patient will begin treatment with:

Azithromycin:

Day 1-3: 500 mg x 1

Day 4-15: 250 mg x 1

If the patient is unable to take the medication orally by themselves, the medication will, if possible, be administered by either stomach-feeding tube, or alternatively, temporary be changed to clarithromycin 500 mg x 2 (this only in agreement with either study coordinator Pradeesh Sivapalan or principal investigator Jens-Ulrik Stæhr Jensen). This will also be done in the control group if necessary. The patient will switch back to azithromycin when possible.

Hydroxychloroquine:

Furthermore, the patient will be treated with hydroxychloroquine as follows:

Day 1-15: 200 mg x 2

(Continued on next page)

(Continued from previous page)

Main outcomes: • Number of days alive and discharged from hospital within 14 days (summarises both whether the patient is alive and discharged from hospital) ("Days alive and out of hospital")

Randomisation: The sponsor (Chronic Obstructive Pulmonary Disease Trial Network, COP:TRIN) generates a randomisation sequence. Randomisation will be in blocks of unknown size and the final allocation will be via an encrypted website (REDCap). There will be stratification for age (>70 years vs. ≤70 years), site of recruitment and whether the patient has any of the following chronic lung diseases: COPD, asthma, bronchiectasis, interstitial lung disease (Yes vs. No).

Blinding (masking): Participants and study personnel will both be blinded, i.e. neither will know which group the participant is allocated to.

Numbers to be randomised (sample size): This study requires 226 patients randomised 1:1 with 113 in each group.

Trial Status: Protocol version 1.8, from April 16, 2020. Recruitment is ongoing (first patient recruited April 6, 2020; final patient expected to be recruited October 31, 2020).

Trial registration: [ClinicalTrials.gov](https://clinicaltrials.gov) Identifier: NCT04322396 (registered March 26, 2020)

Full protocol: The full protocol is attached as an additional file, accessible from the Trials website (Additional file 1). In the interest in expediting dissemination of this material, the familiar formatting has been eliminated; this Letter serves as a summary of the key elements of the full protocol.

The study protocol has been reported in accordance with the Standard Protocol Items: Recommendations for Clinical Interventional Trials (SPIRIT) guidelines (Additional file 2).

Keywords: COVID-19, Randomised controlled trial, protocol, azithromycin, hydroxychloroquine, respiratory infections

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13063-020-04409-9>.

Additional file 1. Full study protocol.

Additional file 2. SPIRIT checklist.

Acknowledgements

We acknowledge all the study personnel involved in ProPAC-COVID.

Authors' contributions

Coordinating investigator - PS; National investigator - JSJ; Design PS, CSU, SBS, PL, CP, TB, TSP, FKK, JDL, TLB, JSJ; Data collection - PS, CSU, RDB, TSL, JVE, JTW, VG, HM, KEJH, CT, JJ, MM, EB, AS, UCSB, DBR, AB, UWM, CBL, JSJ; Analysis PS, JSJ; Writing manuscript PS, CSU, RDB, TSL, JVE, SBS, JTW, VG, PL, HM, KEJH, CT, JJ, MM, EB, AS, CP, UCSB, DBR, AB, UWM, CBL, TB, TSP, FKK, JDL, TLB, JSJ.

Funding

The research project is (investigator) initiated by COP:TRIN. Fund support has been obtained from the Novo Nordisk Foundation of DKK 2,162,646.22 (case no. NNF20SA0062834). The funders had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Availability of data and materials

It is the opinion of the COP:TRIN steering committee that knowledge sharing creates more and better scientific results. Requests for trial information can be submitted to the Project Management (Jens-Ulrik Jensen, Charlotte Ulrik, Pradeesh Sivapalan) who will consider the request. Any reasonable requests will then be discussed with the COP:TRIN Steering Committee.

Ethics approval and consent to participate

The study has been approved by the Danish Committee on Health Research Ethics on April 3, 2020 with ref. no. H-20022574 (for protocol amendment report no. 73409). The study will ensure the protection of human participants according to the Helsinki Declaration and in accordance with Good Clinical Practice Guidelines. Participants (hospital admitted COVID-19 patients) will give written informed consent to participate. A completed patient informed consent form is required from all participants in the study and must be signed by the participant and the informing personnel.

Consent for publication

Not applicable.

Competing interests

PS reports non-financial support from Novartis, personal fees from Boehringer Ingelheim, outside the submitted work. All the other authors declare that they have no competing interests.

Author details

¹Section of Respiratory Medicine, Department of Medicine, Herlev and Gentofte Hospital University of Copenhagen, Hellerup, Denmark. ²Department of Internal Medicine, Zealand University Hospital, Roskilde, Denmark. ³Department of Respiratory Medicine, Amager and Hvidovre Hospital University of Copenhagen, Hvidovre, Denmark. ⁴Department of Surgery, Næstved-Slagelse- Ringsted Hospitals University of Southern Denmark, Slagelse, Denmark. ⁵Department of Respiratory Medicine, Bispebjerg and Frederiksberg Hospital University of Copenhagen, Copenhagen, Denmark. ⁶Department of Respiratory and Infectious Diseases, Nordsjællands Hospital University of Copenhagen, Hillerød, Denmark. ⁷Dep. of Infectious Diseases, Amager and Hvidovre Hospital University of Copenhagen, Hvidovre, Denmark. ⁸Department of Infectious Diseases, Rigshospitalet University of Copenhagen, Copenhagen, Denmark. ⁹Department of Clinical Metabolic Research, Herlev and Gentofte Hospital University of Copenhagen, Hellerup, Denmark. ¹⁰Department of Cardiology,

Herlev and Gentofte Hospital University of Copenhagen, Hellerup, Denmark. ¹¹Department of Respiratory Medicine, Næstved-Slagelse-Ringsted Hospitals University of Southern Denmark, Slagelse, Denmark. ¹²Institute for Regional Health Research University of Southern Denmark, Odense, Denmark. ¹³Department of Respiratory Medicine, Odense University Hospital University of Southern Denmark, Odense, Denmark. ¹⁴Department of Respiratory Medicine, Aalborg University Hospital University of Aalborg, Aalborg, Denmark.

Received: 7 May 2020 Accepted: 14 May 2020

Published online: 10 June 2020

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

