

LETTER

Open Access



BCG revaccination of health workers in Brazil to improve innate immune responses against COVID-19: A structured summary of a study protocol for a randomised controlled trial

Ana Paula Junqueira-Kipnis^{1*} , Laura Raniere Borges dos Anjos¹, Lília Cristina de Souza Barbosa¹, Adeliane Castro da Costa², Kellen Christina Malheiros Borges¹, Amanda da Rocha Oliveira Cardoso³, Kaio Mota Ribeiro¹, Sarah Brena Aparecida Rosa¹, Carine de Castro Souza¹, Rogério Coutinho das Neves¹, Guylherme Saraiva³, Sueli Meira da Silva¹, Erika Aparecida Silveira³, Marcelo Fouad Rabahi³, Marcus Barreto Conte⁴ and André Kipnis¹

Abstract

Objectives: The BCG vaccine, widely used in Brazil in new-borns, induces adjuvant protection for several diseases, including childhood virus infections. BCG activates monocytes and innate memory NK cells which are crucial for the antiviral immune response. Therefore, strategies to prevent COVID-19 in health workers (HW) should be carried out to prevent them becoming unwell so that they can continue to work during the pandemic. The hypothesis is that BCG will improve the innate immune response and prevent symptomatic infection or COVID-19 severity. The primary objective is to verify the effectiveness and safety of the BCG vaccine to prevent or reduce incidence of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection in the city of Goiânia (Brazil) among HW previously vaccinated with BCG and also its severity and mortality during the pandemic of the disease. Secondary objectives are to estimate the incidence of COVID-19 among these professionals and the innate immune response elicited to BCG.

Trial design: This a phase II trial for repositioning BCG as a preventive strategy against COVID-19. The trial is an open-label, parallel-group randomised clinical trial, comparing HW vaccinated with BCG and HW not vaccinated.

Participants: The trial will recruit 800 HW of Goiânia - Goiás, Brazil to reach a total of 400 HW included after comorbidities questioning and laboratorial evaluation.

Eligibility criteria: Any HW presenting BCG vaccination scar with direct contact with suspected COVID-19 patients for (Continued on next page)

* Correspondence: ana_kipnis@ufg.br

¹Institute of Tropical Pathology and Public Health, Federal University of Goiás, Goiânia, Goiás, Brazil

Full list of author information is available at the end of the article



© The Author(s). 2020, corrected publication 2020. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

at least 8 hours per week, whether in hospital beds, ICU, or in transportation or admission (nurses, doctors, physiotherapists, nutritionists, receptionists, etc.) who have negative IgM and IgG COVID-19 test.

Participants with any of the following characteristics will be excluded:

- Have had in the last fifteen days any signs or symptoms of virus infection, including COVID-19;
- Have had fever in the last fifteen days;
- Have been vaccinated fifteen days before the inclusion;
- Have a history or confirmation of any immunosuppressive disease such as HIV, presented solid tumour in the last two years or autoimmune diseases;
- Are under preventive medication with antibiotics, steroid anti-inflammatories, or chemotherapy;
- Have less than 500 neutrophils per mL of blood;
- Have previously been diagnosed with tuberculosis;
- Are breastfeeding or pregnant;
- Are younger than 18 years old;
- Are participating as an investigator in this clinical trial.

Intervention and comparator: HW will be randomized into the BCG vaccinated group or the BCG unvaccinated control group.

The BCG vaccinated group will receive in the right arm, intradermally, a one off dose of 0.1 mL corresponding to approximately 2×10^5 to 8×10^5 CFU of live, freeze-dried, attenuated BCG Moscow 361-I, Bacillus Calmette Guerin vaccine (Serum Institute of India PVT. LTD.).

The unvaccinated control group will not be vaccinated.

The HW allocated in both groups will be followed up at specific times points until 180 days post inclusion.

The vaccinated and control groups will be compared according to COVID-19 related outcomes.

Main outcomes: The primary outcomes are the incidence coefficient of infection by SARS-CoV-2 determined by RT-PCR of naso-oropharyngeal swab specimen or rapid lateral flow IgG and IgM test, and presence of general COVID-19 symptoms, disease severity and admission to hospital during the 180 days of follow up.

The secondary outcome is the innate immune response elicited 15-20 days after vaccination.

Randomisation: The vaccine vial contains approximately 10 doses. In order to optimize the vaccine use, the randomisation was performed in blocks of 20 participants using the platform [randomization.com](http://www.randomization.com) [<http://www.randomization.com>] [<http://www.jerrydallal.com/random/permute.htm>].

The randomization was prepared before any HW inclusion. The results were printed and inserted in sealed envelopes that were numbered with BCG-001 to BCG-400. The printed results as well the envelopes had the same numbers. At the time of the randomisation, each participant that meets the inclusion criteria will receive a consecutive participant number [BCG-001-BCG-400]. The sealed envelope with the assigned number, blinded to the researchers, will be opened in front of the participant and the arm allocation will be known.

Blinding (masking): There is no masking for the participants or for the healthcare providers.

The study will be blinded to the laboratory researchers and to those who will be evaluating the outcomes and performing the statistical analyses. In this case, only the participant identification number will be available.

Numbers to be randomised (sample size): Four hundred health workers will be randomised in two groups. Two hundred participants will be vaccinated, and 200 participants will not be vaccinated.

Trial Status: The protocol approved by the Brazilian Ethical Committee is the seventh version, number CAAE: 31783720.0.0000.5078. The trial has been recruiting since September 20th, 2020. The clinical trial protocol was registered on August 5th, 2020. It is estimated that recruitment will finish by March 2021.

Trial registration: The protocol number was registered on August 5th, 2020 at REBEC (Registro Brasileiro de Ensaios Clínicos). Register number: [RBR-4kjqtg](#) and WHO trial registration number UIN: U1111-1256-3892.

Full protocol: The full protocol is attached as an additional file, accessible from the Trials website (Additional file 1). In the interest in expediting dissemination of this material, the familiar formatting has been eliminated; this Letter serves as a summary of the key elements of the full protocol.

Keywords: COVID-19, Randomised controlled trial, protocol, BCG revaccination, health workers, innate immune response, NK, macrophages, cross-protection

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13063-020-04822-0>.

Additional file 1. Full study protocol

Acknowledgements

The authors thank the collaborators of the project: "Voluntary internship in a randomised clinical trial impacting the COVID-19 pandemic" that included undergraduate, graduate students and professionals that assisted to promote, thru social media, the project BCG-COVID19: Ana Paula dos Santos Rodrigues, Poliana Lopes Parreira, Layanne Mariana Costa da Silva, Ana Carolina de Oliveira Carvalho, Harllon Dayvisson Franco Lucena, Eduarda Pereira Soares Dias, Lays Noleto Nascimento, Luana Michele Alves, Renato Felipe Ferreira Franco, Renato Rodney M. Ferreira, Tayro da Silva Vieira, Thays Noleto Nascimento, Vitor Iago Suriani, Roberta Luiza Rodrigues, Alice Jardim Zaccariotti, Carla Iré Schnier Dias, Daniella Alves Silva, Danielle Dias de Lima, Denise Franco Gontijo dos Santos, Gabriela Santos Silva, Guilherme Vinícius Elias Souza, Guilherme Albuquerque Sampaio, Maria Eduarda Leopoldino da Fonsêca Xavier, Natália Santana Daher Mendonça, Clara Braga dos Santos Azevedo, Raphael Lucas da Silva Marques, Lauren Dalat de Sousa Coelho, Vitória Carreiro de França Teixeira.

The authors also thank Ana Carolina de Oliveira Carvalho for assisting the participant inclusion and Aline de Oliveira Rezende for technical assistance; Laboratory Margarida Dobler Komma from Institute of Tropical Pathology and Public Health - Federal University of Goiás who provided the blood sampling and performed the hemograms of the study participants; Programa Nacional de Imunização for providing the BCG vaccines. The authors are thankful to Dasdores Gonçalves da Silva Oliveira, a certified nurse, that is executing the immunizations of the participants.

Authors' contributions

APJK is the principal investigator, conceived and designed the trial protocol with help from AK, EAS, LRBA, ACC, AROC, GS, MFR and MBC. LRBA and KMR are performing recruitment and study participant scheduling. KCMB, LCSB, CCS, RCN, LRBA are assisting the informed consent collection and participant inclusions. ACC, SBAR, LRBA and KMR are performing the laboratory testing and organizing the sample collection. MBC is supervising the trial statisticians that will be external to the trial study. AROC, GS and MFR are performing the clinical follow up of the study participants for 180 days using telemedicine and/or clinical evaluations. The authors read and approved the final manuscript.

Funding

This trial is funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico/ National Council for Scientific and Technological Development. It received R\$ 1,241,028.00 (approximately US \$206,000.00) that provided equipment and the material costs for the project.

The BCG vaccines were kindly donated by the Brazilian Immunization Program of Health Ministry of the Brazilian Government/ Programa Nacional de Imunização, Ministério da Saúde, Brasil.

The funding agency or the Brazilian Immunization Program has no role in the trial design, conception, collection, results analysis, interpretation or conclusions as well as in the decision to write this manuscript.

Availability of data and materials

The data will be available upon reasonable request to the principal investigator, Dr. Ana Paula Junqueira-Kipnis, ana_kipnis@ufg.br.

Ethics approval and consent to participate

Comissão Nacional de Ética em Pesquisa/ National Research Ethics Commission (CONEP) approved the study. CAAE: 31783720.0.0000.5078. The approval was granted on September 20th, 2020.

Informed consent will be obtained from all participants following a verbal and written explanation of the study, as approved by CONEP.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Institute of Tropical Pathology and Public Health, Federal University of Goiás, Goiânia, Goiás, Brazil. ²Faculdade Estácio de Sá de Goiás – FESGO, Goiânia, Goiás, Brazil. ³Faculty of Medicine, Federal University of Goiás, Goiânia, Goiás, Brazil. ⁴Centro Universitário Arthur Sá Earp Neto. – UNIFASE- Petrópolis, Rio de Janeiro, Brazil.

Received: 30 September 2020 Accepted: 16 October 2020

Published online: 26 October 2020

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

